

AD-A043 376

HARVARD UNIV CAMBRIDGE MA DEPT OF PSYCHOLOGY

F/G 6/16

PREDICTIVE MODELS OF HUMAN VISUAL PROCESSES IN AEROSYSTEMS.(U)

JUL 77 R J MANSFIELD

F44620-76-C-0109


UNCLASSIFIED

AFOSR-TR-77-0977

NL

| OF |
AD
A043 376






NATIONAL BUREAU OF STANDARDS
MICROCOPY RESOLUTION TEST CHART

AD A 043376

12
JES

DDC
RECEIVED
AUG 25 1977
ALBANY NY



Summary

Over the past year a special eye-movement monitoring and visual display system designed to further research on human visual information processing has been partially constructed and is nearing completion. The system can accurately measure eye-movements of less than 1' of arc and with the same precision control the movement of computer-generated visual targets relative to an observer's retina. A section of this report describes the nine subsystems of the visual apparatus and indicates their potential for experimentation. The theoretical reconstruction of the response profile to a visual target is complicated by the fact that under ordinary viewing conditions even during fixation the population of active elements in the visual pathway changes constantly. Using the special visual apparatus, it is possible to maintain a target on a fixed set of retinal receptors facilitating an orderly experimental analysis of visual response patterns. To direct this analysis toward predictive models, theoretical work has been focused on control sites for specific visual functions. At present models of visual transduction in human rods and cones have been devised and several properties numerically evaluated by computer, including temporal modulation transfer functions. Work is currently in progress on a model of visual acuity based upon the spatial modulation transfer function of neurons in the primary visual cortex.

Table of Contents

| | |
|---|----|
| Summary | 1 |
| I. Introduction | 5 |
| II. Equipment | 7 |
| Eyetracker and Image Stabilization System | |
| III. Models | 11 |
| 1. Visual Transduction | |
| 2. Visual Acuity | |
| IV. Projections | 19 |
| V. Bibliography | 21 |

| | |
|---------------------------------|---|
| ACCESSION for | |
| NTIS | White Section <input checked="" type="checkbox"/> |
| DDC | Buff Section <input type="checkbox"/> |
| UNANNOUNCED | <input type="checkbox"/> |
| JUSTIFICATION | |
| BY | |
| DISTRIBUTION/AVAILABILITY NOTES | |
| ON | DATE |
| A | |

I. Introduction

The human visual system is an action-oriented, information-sensing system whose active elements, although grouped into redundant subunits, consist for the most part of several billion interconnecting neurons. The problem of understanding a system of such staggering complexity has been ameliorated by recent scientific advances in the fields of neurobiology and psychophysics and by new technological developments in computation. With present neurobiological techniques it is now possible to determine in precise detail the neural mechanisms subserving perceptual processes and contingent behavioral action (Kandel, 1976). Typically, neurobiological experiments are carried out in non-human animals, but within reasonable limitations the results are applicable to man. In psychophysics, sophisticated methods of linear and quasi-linear analysis have been applied with conspicuous success to detection models for temporal and spatial parameters of visual targets (Kelly, 1972; Robson, 1966; Blakemore and Campbell, 1969). The availability of high-speed computers makes possible the implementation of mathematical models of visual processes too complex for standard analytic solutions. The project takes advantage of all three of these advances.

The approach is to generate mathematical models of visual processes consistent with neurophysiological results, than use psychophysical experiments to estimate parameters and validate the models. Previous quantitative models of human visual processes have been either simple summaries of empirical data of narrow scope with limited predictive value, or theoretical schemas neither in computable form nor realizable in terms

of known or plausible neurophysiological mechanisms (see Bibliography). The novelty of our experimental procedure is to determine the performance characteristics of human vision under conditions of image stabilization. Under conditions of ordinary viewing, even during fixation of a stationary target, the retinal stimulus is never constant due to eye-movements. By initially removing the effects of eye-movements the problem of analyzing and synthesizing the response patterns in the visual pathway is considerably simplified. The system for image stabilization is described in the next section.

II. Equipment

Eyetracker and Image Stabilization System

In order to deconvolve experimentally the contributions of eye-movements from visual processing, we have designed and constructed in part a special optical system. By accurately measuring an observer's eye-movements over a target field, then optically compensating for these movements, the apparatus can effectively stabilize any target on the observer's retina. By the same feedback arrangement, it is possible to superimpose any arbitrary motion of a visual target relative to the observer's retina.

More specifically, the special optical system consists of subsystems as follows:

1. Eye-Movement Monitor: A modified version of the Stanford Research Institute Dual Purkinje Eyetracker (Cornsweet and Crane, 1973; Clark, 1975) measures monocular eye position to an accuracy of 1' of arc with a bandwidth of 150 Hz.
2. Optical Deflector: A custom-designed, high-speed dual optical scanner moves an image plane about the center of rotation of an observer's eye with a repeatability of 1' arc. When appropriately linked to the eye-movement monitor, the total lag time in following eye-movements is at most 1.5 msec, effectively stabilizing the visual target relative to the observer's retina.

3. Image Stabilization Monitor: From the difference in output signals between the eye-movement monitor and the optical deflector, a custom-modified video generator superimposes a marker indicating the observer's point of regard on a video camera image of the target field.
4. Focus Stimulator: A specially-designed lens produces changes in optical focus without changing target size or brightness so that the system has the capability of moving targets in three dimensions.
5. Visual Display: Visual targets can be presented by means of three display devices: a CRT, a Maxwellian view stimulator, and a Video monitor.
 - (a) The CRT display can present two multiplexed channels of luminous bars or sinusoidally-varying luminance gratings whose extent, position, and contrast can be changed at rates up to 1000 Hz.
 - (b) The Maxwellian optical system can provide up to three channels of controlled target or background illumination of varied spectral composition over a wide range of intensities up to 10^7 for white light.
 - (c) The Video display can present patterns of 512 by 512 elements with a 6-bit gray scale.

6. Data Acquisition and Control Interface: For each of the above devices, computer-control and data sampling are available in flexible form through an interconnection panel. The interface contains 16 TTL input lines, 16 TTL output lines, 16 multiplexed 12-bit A/D input lines, 8 12-bit D/A output lines, and a DMA channel for the video generator.
7. Laboratory Computer: An Eclipse S/200 with 32 K words and 10 Mbyte disc is available to generate visual displays, control experiments, collect and analyze experimental data.




Figure 1. Theoretical impulse response functions for a human rod photoreceptor. For each energy level, given in relative units, the time-course of membrane hyperpolarization has been calculated from a mathematical model of visual transduction. With increasing pulse energy, rise-time decreases and peak amplitude saturates.

III. Models

The first year of the project has been devoted mainly to the generation of computer software and the construction of apparatus; however, some preliminary models have been devised for later experimental testing. Our strategy has been to focus attention on critical control points in visual processing. For example, extensive neurophysiological evidence supports the view that the only essential non-linearity associated with intensity coding in vision and many other senses is confined to the periphery (Mountcastle, 1967; Stevens, 1970). Modelling the nonlinear transformation in the periphery sets the stage for the application of linear models for central processing.

1. Visual Transduction (with J. Daugman)

Electrical recordings from vertebrate retinas over the past decade have thoroughly discredited the photocell theory of visual transduction. It is now clear that vertebrate photoreceptors are active elements with complex response properties that to a considerable extent quantitatively account for the temporal and intensive discrimination capacity of the intact visual system (e.g., Boynton and Whitten, 1970; Kelly, Boynton, and Baron, 1976). Individual primate rod and cone photoreceptors have not been recorded for the prolonged periods necessary to obtain extensive quantitative results, but the initial findings suggest a close qualitative similarity to other vertebrate photoreceptors (DeMonasterio and Gouras, personal communication). So, in devising mathematical models for human rod and cone photoreceptors we have

been guided by the work of others on photoreceptors in non-mammalian vertebrates (Baylor, Hodgkin, and Lamb, 1974; Cervetto, Pisano, and Torres, 1977), taking into account the particular dimensions of human photoreceptors and the higher operating temperature (37.5° C).

Figure 1 shows the calculated response for a human rod photoreceptor to a series of light impulses of varying energy content. Two types of nonlinearity are evident in the response. First, a saturating nonlinearity for, as the energy is increased in equal ratio steps, the peak amplitude increases linearly at low levels, but approaches an asymptotic value at high levels. To a first approximation, the amplitude of the peak, V , can be described as a function of flash energy, E , by the following equation:

$$\frac{V}{V_{\max}} = \frac{E}{E + s_1} \quad (1)$$

where V_{\max} is the maximum attainable response, and s_1 is a semi-saturation constant. Equation (1) has been found to describe the amplitude of response to brief flashes in several species of vertebrates (Mansfield, 1976). The second type of nonlinearity found is a time-scale nonlinearity. As the energy increases, the rise-time remains constant at low levels, then decreases to an asymptotic value at high levels. To a first approximation, the rise-time of the peak, t , can be described as a function of flash energy, E , by the following equation:




Figure 2. Theoretical transfer functions for a human foveal cone. Amplitude sensitivity for three background levels of quantum flux is plotted as a function of temporal frequency of modulation based upon calculations from a model of visual transduction. Rather than being of constant shape as in a linear system, the shape of the transfer function is level-dependent.

$$\frac{t}{t_{\min}} = \left[\frac{I^{1/3}}{I^{1/3} + s_2} \right]^{-1} \quad (2)$$

where t_{\min} is the minimum attainable rise-time and s_2 is a constant. Equation (2) has been found to describe the latency of response to brief flashes in a number of vertebrate species (Mansfield, 1973; Mansfield and Daugman, 1977). Despite these two nonlinearities found in the response characteristics of the model for human photoreceptors and those measured electrophysiologically in other vertebrates, a steady-state condition of constant illumination produces a linearization of photoreceptor response. Using the resulting small signal linearity prevailing at each level of light intensity, transfer functions can be calculated.

Figure 2 shows the amplitude component of transfer functions calculated for a human foveal cone for three light intensity levels. For clarity of presentation and for comparison with human psychophysical data for flicker thresholds, the functions are plotted using an ordinate in units of sensitivity (nanovolts/photon/cone-second). The shape of the family of curves resembles that obtained by Kelly (1971) for counterphase gratings which eliminate the familiar low-frequency attenuation by reducing lateral interactions in the proximal portions of the retina beyond the photoreceptors. Such a result is encouraging for it suggests that by obtaining the product of the transfer function for the appropriate portion of the proximal inhibitory network and the transfer function for the

appropriate type of photoreceptor, it will be possible to predict the form of the human detection threshold curve for a wide variety of spatio-temporal patterns.

2. Visual Acuity (with S. Ronner and G. Legge)

The threshold contrast for detecting a visual target with a particular spatial configuration is influenced by a number of factors, but under conditions where the nonlinear transformations in the retina can be set aside so that the response of the visual system can be considered linear, the cortical mosaic of neurons with spatially complex receptive fields functions as a major control site. At the level of the receptive field structure of striate cortical neurons, the spatial inhomogeneity and anisotropy of primate vision become clearly evident (Hubel and Wiesel, 1974; Mansfield, 1974). Fortunately, the inhomogeneity and anisotropy are orderly in nature. The average size of receptive fields is small in the foveal projection region of primary visual cortex, but increases approximately as the distance from the fovea. In addition, receptive fields preferentially sensitive to horizontal and vertical targets predominate in the foveal projection region, but not in the periphery. Were it not for the inhomogeneity and anisotropy a predictive model could be constructed from a single linear spatial filter using the Fourier transform of the unique point spread function.

As an instance of the detection problem we have considered the case of oriented line segments. To construct an appropriate spatial filter that is consistent with neurophysiological data,




Figure 3. Orientation selectivity in primate vision. The solid squares represent the threshold elevation in human observers following adaptation to a high contrast sine wave grating. The open circles represent the integrated impulse discharge elicited by a luminous narrow bar in a monkey visual neuron recorded in the foveal projection region of Area 17. The solid line fitted to the neural data represents the orientation response profile calculated for a theoretical spatial filter.

subunits, each of which has an identical spatial impulse response function of the form

$$F(r) = e^{-r^2} - 0.04 e^{-r^2/25}, \quad (3)$$

are combined in arrays and their outputs summed linearly. The response profile of such subunits resembles that found for sustained on-center off-surround mammalian visual neurons at the level of the retinal ganglion cells or lateral geniculate. For example, in the cat lateral geniculate the radius of the surround area is typically five times larger than the radius of the center area (Enroth-Cugell and Robson, 1966). When a line segment function is convoluted with the spatial filter at different relative orientations, an orientation response function is obtained such as that shown in Figure 3.

Figure 3 shows an orientation response function based upon nine subunits and possessing a half-maximal bandwidth of 48° . For purposes of comparison, two sets of data are shown: orientation sensitivity measures for a human observer derived psychophysically for a sine-wave grating target, and orientation response profile for a monkey visual neuron in the foveal projection region of Area 17. Both sets of data are in accord and are well fitted by the theoretical curve generated by the model. Since the bandwidth of the neuron is close to the mode of the population (Ronner, Legge, and Mansfield, 1976), the orientation sensitivity of the most numerous neurons may well be the determinant of threshold for the psychophysical task. By incorporating distribution parameters for

the spatial filters constructed from a cluster of subunits, it should be possible to account for inhomogeneity and anisotropy across the visual field and devise a more general model for the detection of visual targets.

IV. Projections

The understanding of vision in biological systems in terms of predictive mathematical models is a central problem from the scientific point of view whose solution is of great practical significance. How does the human visual system recognize patterns or reconstruct the visual world? Is a Fourier-like decomposition an intrinsic part of the process (Ginsburg, 1973)? At present our research has taken some initial steps toward answering these intriguing questions. Dynamic models for human rod and cone photoreceptors have been generated and the effects of cortical organization on the threshold detection of stationary, oriented targets examined. These models will serve as building blocks in the synthesis of more complex visual processes. The next step is to evaluate the models in psychophysical experiments and incorporate into more general models the effects of retinal adaptation and inhibitory spatial interactions. Such effects need to be studied in the absence of eye-movements. By stabilizing the images of visual targets on the observer's retina, it will be possible to obtain precise estimates of the parameters necessary to evaluate and extend the models.

V. Bibliography

- Baylor, D. A., and Fuortes, M. G. F. (1970). Electrical responses of single cones in the retina of the turtle. J. Physiol., 207, 77-92.
- Baylor, D. A., Hodgkin, A. L., and Lamb, T. D. (1974). Reconstruction of the electrical responses of turtle cones to flashes and steps of light. J. Physiol., 242, 759-791.
- Blakemore, C., and Campbell, F. W. (1969). On the existence of neurons in the human visual system selectively sensitive to the orientation and size of retinal images. J. Physiol., 203, 237-260.
- Blakemore, C., Nachmias, J., and Sutton, P. (1970). The perceived spatial frequency shift: evidence for frequency selective neurones in the human brain. J. Physiol., 210, 727-750.
- Boynton, R. M., and Whitten, D. N. (1970). Visual adaptation in monkey cones: recordings of late receptor potentials. Science, 170, 1423-1426.
- Campbell, F. W., Carpenter, R. H. S., and Levinson, J. Z. (1969). Visibility of aperiodic patterns compared with that of sinusoidal gratings. J. Physiol., 204, 283-298.
- Campbell, F. W., and Green, D. G. (1965). Optical and retinal factors affecting visual resolution. J. Physiol., 181, 576-593.
- Campbell, F. W., and Gubish, R. W. (1966). Optical quality of the human eye. J. Physiol., 186, 558-578.
- Campbell, F. W., Howell, E. R., and Robson, J. G. (1971). The appearance of gratings with and without the fundamental Fourier component. J. Physiol., 217, 16P-17P.

- Campbell, F. W., and Kulikowski, J. J. (1966). Orientational selectivity of the human visual system. J. Physiol., 187, 437-445.
- Campbell, F. W., Kulikowski, J. J., and Levinson, J. (1966). The effect of orientation on the visual resolution of gratings. J. Physiol., 187, 427-436.
- Campbell, F. W., and Maffei, L. (1970). Electrophysiological evidence for the existence of orientation and size detectors in the human visual system. J. Physiol., 207, 635-652.
- Cervetto, L., Pisano, E., and Torres, V. (1977). Electrical responses of rods in the retina of Bufo marinus. J. Physiol., 267, 17-51.
- Childress, D. S., and Jones, R. W. (1967). Mechanics of horizontal movement of the human eye. J. Physiol., 188, 273-284.
- Clark, M. R. (1975). A two-dimensional Purkinje eye tracker. Behavior Research Methods and Instrumentation, 7, 215-219.
- Cornsweet, T. N., and Crane, H. D. (1973). Accurate two-dimensional eye tracker using first and fourth Purkinje images. J. Opt. Soc. Amer., 63, 921-928.
- DeValois, R. L. (1965). Behavioral and electrophysiological studies in the primate visual system. In: W. D. Neff (Ed.), Contributions to Sensory Physiology, New York, Academic Press, 1, pp. 137-178.
- Ditchburn, R. W. (1973). Eye-movements and Visual Perception. Oxford, Clarendon Press.
- Dowling, J. E. (1967). The site of visual adaptation. Science, 155, 273-279.

- Dowling, J. E., and Ripps, H. (1970). Visual adaptation in the retina of the skate. J. Gen. Physiol., 56, 491-520.
- Enroth-Cugell, C., and Robson, J. G. (1966). The contrast sensitivity of retinal ganglion cells in the cat. J. Physiol., 187, 517-552.
- Fain, G. L., and Dowling, J. E. (1973). Intracellular recordings from single rods and cones in the mudpuppy retina. Science, 180, 1178-1181.
- Ginsburg, A. P. (1973). Pattern recognition techniques suggested from psychological correlates of a model of the human visual system. Proceedings IEEE 1973 NAECON, Dayton, Ohio, pp. 309-316.
- Hines, M. (1976). Line spread function variation near the fovea. Vision Res., 16, 567-572.
- Hubel, D. H., and Wiesel, T. N. (1974). Uniformity of monkey striate cortex: a parallel relationship between field size, scatter, and magnification factors. J. Comp. Neurology, 158, 295-306.
- Kandel, E. R. (1976). Cellular Basis of Behavior. San Francisco, W. H. Freeman.
- Kelly, D. H. (1971). Theory of flicker and transient responses. II. Counterphase gratings. J. Opt. Soc. Amer., 61, 632-640.
- Kelly, D. H. (1972). Adaptation effects on spatio-temporal sine-wave thresholds. Vision Res., 12, 89-101.
- Kelly, D. H., Boynton, R. M., and Baron, W. S. (1976). Primate flicker sensitivity: psychophysics and electrophysiology. Science, 194, 1077-1079.
- Land, E. H. (1959a). Color vision and the natural image. Part I. Proc. Natl. Acad. Sci., 45, 115-129.

- Land, E. H. (1959b). Color vision and the natural image. Part II. Proc. Natl. Acad. Sci., 45, 636-644.
- Land, E. H. (1964). The retinex. Amer. Scientist, 52, 247-264.
- Land, E. H., and McCann, J. J. (1971). Lightness and retinex theory. J. Opt. Soc. Amer., 61, 1-11.
- Legge, G. E. (1976). Contrast detection in human vision: spatial and temporal properties. Doctoral dissertation, Harvard University, Cambridge, Massachusetts.
- Macleod, I. D. G., and Rosenfeld, A. (1974). The visibility of gratings: spatial frequency channels or bar detecting units. Vision Res., 14, 909-915.
- Mansfield, R. J. W. (1973a). Brightness function: effect of area and duration. J. Opt. Soc. Amer., 63, 913-920.
- Mansfield, R. J. W. (1973b). Latency functions in human vision. Vision Res., 13, 2219-2234.
- Mansfield, R. J. W. (1974). Measurement, invariance, and psychophysics. In: H. R. Moskowitz, B. Scharf, and J. C. Stevens (Eds.), Sensation and Measurement - Papers in Honor of S. S. Stevens, Dordrecht, Netherlands, Reidel Press, pp. 111-126.
- Mansfield, R. J. W. (1976). Visual adaptation: retinal transduction, brightness, and sensitivity. Vision Res., 16, 679-690.
- Mansfield, R. J. W., and Daugman, J. G. (1977). Retinal mechanisms of visual latency. Submitted for publication.
- Mansfield, R. J. W., Poggio, G. F., and Sillito, A. (1977). Neural mechanisms of color vision: spectral sensitivity and hue discrimination. In preparation.

- Mountcastle, V. B. (1967). The problem of sensing and the neural coding of sensory events. In: G. C. Quarten, T. Melnechuk, and F. C. Schmitt (Eds.), The Neurosciences, New York, Rockefeller University Press, pp. 393-408.
- Norman, R. A., and Werblin, F. S. (1974). Control of retinal sensitivity I. Light and dark adaptation of vertebrate rods and cones. J. Gen. Physiol., 63, 37-61.
- Poggio, G. F., Baker, F. H., Mansfield, R. J. W., Sillito, A., and Grigg, P. (1975). Spatial and chromatic properties of neurons subserving foveal and parafoveal vision in Rhesus monkey. Brain Res., 100, 25-59.
- Robinson, D. A. (1964). The mechanics of human saccadic eye movement. J. Physiol., 174, 245-264.
- Robinson, D. A. (1968). Eye movement control in primates. Science, 161, 1219-1224.
- Robson, J. G. (1966). Spatial and temporal contrast-sensitivity functions of the visual system. J. Opt. Soc. Amer., 56, 1141-1142.
- Ronner, S. F., Legge, G. E., and Mansfield, R. J. W. (1976). Neural mechanisms of orientation discrimination in primate vision. Paper presented at the Spring meeting of the Association for Research in Vision and Ophthalmology, Sarasota, Florida.
- Rushton, W. A. H. (1965). The Ferrier Lecture, 1962: Visual adaptation. Proc. Roy. Soc. (London), ser. B, 162, 20-46.
- Skavenski, A. (1972). Inflow as a source of extra-retinal eye position information. Vision Res., 12, 225-236.

- Sperling, G., and Sondhi, M. M. (1968). Model for visual luminance discrimination and flicker detection. J. Opt. Soc. Amer., 58, 1133-1145.
- Stevens, S. S. (1970). Neural events and the psychophysical law. Science, 170, 1043-1050.
- Stockham, T. G., Jr. (1972a). Intraframe encoding for monochrome images by means of a psychophysical model based on nonlinear filtering of multiplied signals. In: T. S. Huang and O. J. Tretiak (Eds.), Proceedings 1969 Symposium Picture Bandwidth Compression, New York, Gordon and Breach, pp. 415-442.
- Stockham, T. G., Jr. (1972b). Image processing in the context of a visual model. Proceedings IEEE, 60, 828-842.
- Thomas, J. P. (1970). Model of the function of receptive fields in human vision. Psychol. Rev., 77, 121-134.
- Yarbus, A. L. (1967). Eye Movements and Vision. New York, Plenum Press.

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|--|-----------------------|---|
| 1. REPORT NUMBER (6) AFOSR-TR-77-0977 | 2. GOVT ACCESSION NO. | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) PREDICTIVE MODELS OF HUMAN VISUAL PROCESSES IN AEROSYSTEMS. | | 5. TYPE OF REPORT & PERIOD COVERED Scientific Interim Report 1 June 1976 to 31 May 1977 |
| | | 6. PERFORMING ORG. REPORT NUMBER |
| 7. AUTHOR(s) (10) Richard J. W. Mansfield | (15) | 8. CONTRACT OR GRANT NUMBER(s) F44620-76-C-0109 |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS Harvard University 1350 Massachusetts Avenue Cambridge, Massachusetts 02138 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F/2313A4 (12) A4 |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research (NL) Building 410 Bolling Air Force Base, D. C. 20332 | | 12. REPORT DATE July, 1977 |
| | | 13. NUMBER OF PAGES 26 |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) (7) Annual rept. 1 Jun 76 - 31 May 77 | | 15. SECURITY CLASS. (of this report) Unclassified |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited | | 15a. DECLASSIFICATION DOWNGRADING SCHEDULE (12) 26p. |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Vision, Mathematical Models, Linear Systems Analysis, Eye-Movement Monitoring, Computer Generated Visual Displays. | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Over the past year a special eye-movement monitoring and visual display system designed to further research on human visual information processing has been partially constructed and is nearing completion. The system can accurately measure eye-movements of less than 1' of arc and with the same | | |

→ precision control the movement of computer-generated visual targets relative to an observer's retina. A section of this report describes the 9 subsystems of the visual apparatus and indicates their potential for experimentation. The theoretical reconstruction of the response profile to a visual target is complicated by the fact that under ordinary viewing conditions even during fixation the population of active elements in the visual pathway changes constantly. Using the special visual apparatus, it is possible to maintain a target on a fixed set of retinal receptors facilitating an orderly experimental analysis of visual response patterns. → To direct this analysis toward predictive models, theoretical work has been focused on control sites for specific visual functions. At present, models of visual transduction in human rods and cones have been devised and several properties numerically evaluated by computer, including temporal modulation transfer functions. Work is currently in progress on a model of visual acuity based on the spatial modulation transfer function of neurons in the primary visual cortex.

Unclassified